## **Assignment**

#### **LESSON 2: Bow Thai**

#### Write

Determine whether each statement is true or false. If the statement is false, rewrite it so that it becomes a true statement.


- 1. A translation function along parallel lines is a type of geometric transformation that is non-rigid.
- 2. Isometries include geometric transformations such as translations, rotations, and reflections.
- 3. A transformation is an isometry if it does not preserve size and shape.
- 4. A dilation is a non-rigid geometric transformation.

#### Remember

A translation is a function, T, which takes as its input a set of pre-image points and outputs a set of image points. The pre-image points are translated a distance of AB in the direction AB. For example, a translation of point P could be expressed as  $T_{AB}(P)$ , or P'.

#### **Practice**


1. Write a function to describe each translation. Each solid figure is a pre-image, and each dashed figure is an image.







b.  $T_{I'M}$ (Triangle) = Triangle'




- 2. Complete each translation given the function.
- a.  $T_{AB}$ (Rectangle) = Rectangle'









LESSON 2: Bow Thai • 145


# **Assignment**

#### Write


- 1. False; A translation function along parallel lines is a type of geometric transformation that is rigid.
- 2. True
- 3. False: A transformation is an isometry if it does preserve size and shape.
- 4. True

#### **Practice**


- 1a. Sample answer.  $T_{AB}$  (Parallelogram) = Parallelogram'
- 1b. Sample answer.  $T_{BA}$  (Trapezoid) = Trapezoid'
- 1c. Sample answer.  $T_{PO}$  (Hexagon) = Hexagon'
- 1d. Sample answer.  $T_{OP}$  (Triangle) = Triangle'
- 2a.




2b.








2d.



#### Assignment Answers


#### **Stretch**




1b. Each point of the trapezoid moved a total distance of  $\sqrt{85}$  along a line with a slope of  $\frac{2}{9}$ .

### **Review**

1.



2.



#### Stretch

- 1. A translation function for a trapezoid is given as  $T_{pQ}$  (Trapezoid ABCD) = Trapezoid A'B'C'D'. Trapezoid ABCD has vertices A (-7, -1), B (-5, -1), C (-4, -4), and D (-8, -4). If P is located at (-2, 3) and Q is located at (7, 5), determine the vertices of the translated trapezoid A'B'C'D'.
  - a. Draw points *P* and *Q* and trapezoids *ABCD* and *A'B'C'D'* on a coordinate plane. Include a dashed line between points *P* and *Q*.
  - b. Determine the distance traveled by each point of the trapezoid and the slope of the line along which the points moved.

#### Review

- 1. Construct a square inscribed in a circle using the given line segment.
- 2. Construct a regular hexagon inscribed inside circle A.



- 3. Identify the mappings that are functions. If the mapping is not a function, explain why not.
  - a. {(1, 1), (1, 3), (3, 1), (2, 2), (5, 5)}
  - b. {(2, 4), (3, 5), (4, 6), (6, 8), (8, 10)}
  - c. {(-1, 2), (0, 1), (1, 1), (2, 1), (3, 1), (4, 0)}

146 • TOPIC 2: Rigid Motions on a Plane

- 3a. This mapping is not a function. A single x-value (1) is mapped to two different y-values (1 and 3).
- 3b. This mapping is a function.
- 3c. This mapping is a function.